
A revised checklist of the French Culicidae

Francis Schaffner

17, rue du 152e R.I. – 68700 Steinbach, France

There are few major works on French mosquitoes. Those that exist are species catalogues (Séguy, 1925a; Hedeon, 1958, 1959; Callot & Rioux, 1965), sometimes including distribution and bibliography (Rageau *et al.*, 1970; Moussiegt, 1986). A certain number of species listed in the literature cannot be considered as truly belonging to the French fauna. Furthermore, during the last few years, new species have been described, and taxonomic changes have occurred (synonyms, new sub-genera, etc.). The purpose of this paper is to update the list of species collected in France and Corsica.

Sub-family ANOPHELINAE

Genus *Anopheles* Meigen 1818

Sub-genus *Anopheles* Meigen 1818

- 1 *algeriensis* Theobald 1903
- claviger sensu lato*:
- 2 *claviger* (Meigen 1804)
- 3* *petragnanii* Del Vecchio 1939
- 4 *hyrcanus* (Pallas 1771)
- maculipennis sensu lato*:
- 5 *atroparvus* Van Thiel 1927
- 6** *labranchiae* Falleroni 1926
- 7 *maculipennis s.s.* Meigen 1818
- 8 *melanoon* Hackett 1934
- 9 *messeae* Falleroni 1926
- 10** *sacharovi* Favre 1903
- 11 *subalpinus* Hackett & Lewis 1935
- 12** *marteri* Senevet & Prunelle 1927
- 13 *plumbeus* Stephens 1828

Sub-genus *Cellia* Theobald 1902

- 14** *superpictus* Grassi 1899

Sub-family CULICINAE

Tribe *Aedini*

Genus *Aedes* Meigen 1818

Sub-genus *Aedes* Meigen 1818

- 15 *cinereus* Meigen 1818
- 16 *esoensis rossicus* Dolbeskin, Gorickaja & Mitrofanova 1930
- 17 *geminus* Peus 1970

Sub-genus *Aedimorphus* Theobald 1903

- 18 *vexans vexans* (Meigen 1830)
- 19 *vittatus* (Bigot 1861)

Sub-genus *Finlaya* Theobald 1903

- 20 *geniculatus* (Olivier 1791)

Sub-genus *Ochlerotatus* Lynch Arribalzaga 1891

Intrudens Group

- 21 *diantaeus* Howard, Dyar & Knab 1912
- 22 *pullatus* (Coquillett 1904)

Ochlerotatus Group

Annulipes Sub-group

- 23 *annulipes* (Meigen 1830)
- 24 *cantans* (Meigen 1818)
- 25 *flavescens* (Müller 1764)
- 26 *surcoufi* Theobald 1912

Communis Sub-group

- 27 *cataphylla* Dyar 1916
- 28 *communis* (De Geer 1776)
- detritus* (Haliday 1833)
- 29 *sp. A* Pasteur *et al.* 1977
- 30 *sp. B* Pasteur *et al.* 1977
- 31 *nigrinus* (Eckstein 1918)
- 32 *punctor* (Kirby 1837)
- 33 *sticticus* (Meigen 1838)

Dorsalis Sub-group

- 34 *berlandi* Séguy 1921
- caspius caspius* (Pallas 1771)
- 35 *sp. A* Cianchi *et al.* 1980
- 36 *dorsalis* (Meigen 1830)
- 37 *mariae* (Sergent & Sergent 1903)
- 38 *pulcritarsis pulcritarsis* (Rondani 1872)

Sub-genus *Rusticoides* Shevchenko & Prudkina 1973

- 39 *refiki* Medschid 1928
- 40 *rusticus* (Rossi 1790)

Sub-genus *Stegomyia*

- 41*** *aegypti* (Linnaeus 1762)

Tribe *Culicini*

Genus *Culex* Linnaeus 1758

Sub-genus *Barraudius* Edwards 1921

- 42 *modestus modestus* Ficalbi 1889

Sub-genus *Culex* Linnaeus 1758

- 43** *brumpti* Galliard 1931
- 44 *mimeticus* Noe 1899
- 45 *pipiens pipiens* Linnaeus 1758 inc. form *molestus* Forskål 1775
- 46 *theileri* Theobald, 1903
- 47 *torrentium* Martini 1925

Sub-genus *Maillotia* Theobald 1907

- 48 *hortensis hortensis* Ficalbi 1889

Sub-genus *Neoculex* Dyar 1905

- 49 *impudicus* Ficalbi 1890
- 50 *martinii* Medschid 1930
- 51 *territans* Walker 1856

Tribe *Culisetini*

Genus *Culiseta* Felt, 1904

Sub-genus *Allotheobaldia* Brolemann 1919

52 *longiareolata* (Macquart 1838)

Sub-genus *Culicella* Felt 1904

53 *fumipennis* (Stephens 1825)

54 *litorea* (Shute 1928)

55 *morsitans* (Theobald 1901)

Sub-genus *Culiseta* Felt 1904

56 *alaskaensis alaskaensis* (Ludlow 1906)

57 *annulata* (Schränk 1776)

58 *glaphyoptera* (Schiner 1864)

59 *subochrea* (Edwards 1921)

Tribe *Mansoniini*

Genus *Coquillettidia* Dyar 1905

Sub-genus *Coquillettidia* Dyar 1905

60 *buxtoni* (Edwards 1923)

61 *richiardi* (Ficalbi 1889)

Tribe *Orthopodomyiini*

Genus *Orthopodomyia* Theobald 1904

62 *pulcripalpis* (Rondani 1872)

Tribe *Uranotaeniini*

Genus *Uranotaenia* Lynch Arribalzaga 1891

Sub-genus *Pseudoficalbia* Theobald 1912

63 *unguiculata unguiculata* Edwards 1913

* this taxon should be *petragnanii* and not *petragnani* following the Latin rules of declension for specific names. The species was named by Del Vecchio in honour of Pr. Petragnani

** species present in Corsica only

*** sporadic species, often introduced via ports, not regarded as endemic in view of the rarity of observations (Rageau *et al.*, 1970)

COMMENTS

I. Arrangement of the taxonomic list

The organisation of this list takes into account the systematic classification used in the reference work: "A catalog of the mosquitoes of the world" and its additions (Knight & Stone, 1977; Knight, 1978; Ward, 1984; Gaffigan & Ward, 1985; Ward, 1992). Taxa are presented alphabetically within each taxonomic grouping.

Harbach (1994) has recently revised the internal classification of the genus *Anopheles*. Considering the small number of species, it was decided not to represent the sections, series, groups and sub-groups. On the other hand, certain relationships are deliberately highlighted by reference to "species complexes".

Considering that the sub-genus *Ochlerotatus* represents by far the greatest number of species, and that the morphological differences are sometimes important, it was decided to arrange the species according to the groups and sub-groups used by Mohrig (1969).

The sub-genus *Rusticoides* was described by Shevchenko & Prudkina (1973) following a study of the male genitalia in the genus *Aedes*. They proposed *Ae. refiki* as haplotype, and included in the sub-genus the related species previously incorporated in the *rusticus* group. Even though the catalogue of mosquitoes (Knight & Stone, 1977 and its additions) classifies *Ae. refiki* as being the only European species in this sub-genus, we share the opinion of other authors (Shevchenko & Prudkina, 1973; Encinas Grandes, 1982) that *Ae. rusticus* is also a member of that sub-genus.

II. The Species list

We have made several modifications to the last list of French Culicidae published by Callot & Rioux (1965). Their list recognised 62 taxa, including *Ae. vexans* which was accidentally omitted from the publication due to a typographical error.

Species added to the list

An. subalpinus is a taxon that has recently been elevated to species status (Bullini *et al.*, 1980; Cianchi *et al.*, 1987; Ribeiro *et al.*, 1988; Ward, 1992). Being morphologically very close to *An. melanoon*, the two species are often mistaken in the literature. *An. subalpinus* was first recorded in the Languedoc in 1957 (Rioux & Ruffié, 1957) and appears to be abundant in the south of France (Salieres *et al.*, 1978).

Ae. esoensis rossicus is the most recently discovered species in France (Schaffner & Pfirsch, 1995). It has been recorded only in the north of Alsace, where it is relatively rare. However, large populations of this species occur in the flood plains of the Rhine in Germany (Becker & Ludwig, 1981).

Ae. geminus is a species related to *Ae. cinereus*, as described by Peus (1970). Since its discovery in France (Roman & Pichot, 1976), this species has been only rarely recorded, probably due to the difficulties posed by its identification (the need to distinguish adult males). It seems, however, that *Ae. geminus* is more abundant than *Ae. cinereus* (Schaffner, 1992; Schaffner & Pfirsch, 1995). Besides, it seems likely that a certain number of earlier records of *Ae. cinereus* are actually attributable to *Ae. geminus*. The distributions of both these taxa are still to be established.

Sibling species have been differentiated within two taxa of the genus *Aedes* (*Ae. detritus* and *Ae. caspius caspius*) using new genetic identification techniques, in particular isoenzymes studies. In addition to the demonstrated reproductive isolation, ecological, ethological and distributional differentiation has been added, thus confirming the validity of the two *Ae. detritus* species.

- *Ae. detritus* (both species "A" and "B" of Pasteur *et al.* (1977)), were recorded in Camargue (Pasteur *et al.*, 1977). A geographical segregation has been found to exist along a north-south gradient, with "A" less common in the north (Atlantic coastline: Brutus *et al.*, 1994). Autogeny, frequent in "A", is exceptional in "B"; species "A" develops only in hyper-saline larval sites; and the possibility of finding some morphological differentiation characters cannot be excluded (Agoulon, 1996).
- *Ae. caspius caspius*: only species "A" of Cianchi *et al.*, 1980 has been identified in France, so far, on the Atlantic coastline (Agoulon, 1996).

A change in nomenclature

Arnaud *et al.* (1976) resurrected the taxon *Aedes surcoufi* from synonymy with *Aedes (Ochlerotatus) excrucians* (Walker 1856) and considered it to be the sole representative of the "excrucians" complex in France. Accordingly, the name *Ae. excrucians* is replaced by *Ae. surcoufi* in the list of species. Certain authors (Dahl, C. & Nielsen, B.O. personal communication) consider, however, that the possibility of finding *Ae. excrucians* in the Alps should not be excluded. In the absence of any morphological analysis of alpine populations, we hold the opinion of Arnaud *et al.* (1976).

Species removed from the list

Aedes (Ochlerotatus) leucomelas (Meigen 1804) is removed from the list since it is not found in France. The only reports noted the occurrence of aggressive *Ae. leucomelas* females in the Jura (Roman, 1944; Roman & Morel, 1946), but emphasised the difficulties of distinguishing this species from *Aedes cataphylla*. A few years later the same specimens were classified as *Ae. cataphylla* (Roman, 1958). Hedeën (1959) and Rageau *et al.* (1970) subsequently suggested that the species probably occurred in France, considering its extensive Palaearctic range. However, as subsequent studies failed to locate the species in France, it seems reasonable to exclude it from the list. Nevertheless, *Ae. leucomelas* occurs in Germany, a few kilometres from the French border (Oberrheingebiet) (Becker & Ludwig, 1981) and therefore could potentially be encountered in the north-east of France.

Culex (Culex) laticinctus Edwards 1913 has been recorded in three different places: Hyères, in Var (Séguy, 1920a), Fontainebleau, in Seine-et-Marne (Séguy, 1921) and Lucciana, in Corsica (Galliard, 1927). All other references to this species in France actually relate to one or more of these records. We consider that the presence of a species can only be authenticated after larvae, adult males and adult females have been observed, together with the possibility of studying preserved specimens. None of the three French records matches these requirements. In addition to this is the fact that the species distribution is disjunct (Fontainebleau). We therefore do not consider this species to be present in France, at least until there are further records and more studies have been made.

Culex (Culex) univittatus Theobald 1901 was recorded in Corsica (Galliard, 1927) on the basis of larvae attributed to *Culex perexiguus* Theobald, 1903. Subsequently, the same author identified adult females from two other Corsican sites, but mentioned that there was no certainty of these specimens being *Cx. univittatus* (= *perexiguus*) as he had failed to obtain adults from the previously mentioned larvae (Galliard, 1928). Later, the discovery in 1931 of similar larvae from one of the 1927 sites, and the obtaining of adults (2 males and 2 females) allowed the author to describe the specimens as belonging to the new taxon *Culex brumpti* Galliard 1931. As no precise detail was given for the females from 1928, they remained classified as *Cx. univittatus*. However, even though various authors have retained them in their faunistic lists (Edwards, 1928; Séguy, 1932; Callot & Rioux, 1965; Moussiegt, 1986), others have removed the taxon (Lanzalavi, 1965; Rageau *et al.*, 1970). In 1966, there was another mention of the species in an administrative document from Corsica (Ségui, 1966). The species was encountered three times, with a total of 4 larvae (no preserved specimens). Even though *Cx. univittatus* occurs in the Iberian Peninsula and in Italy (Dahl & White, 1978), we prefer not to include it in the French species list, due to the uncertainties in the literature.

III. Other species and sub-species not included

Apart from the list by Callot & Rioux (1965) and species recently reported, the literature contains records of five other taxa in France. Four of these are invalid and one is considered doubtful.

Aedes (Finlaya) echinus (Edwards 1920) was reported for the first time (Séguy, 1920b) after a female specimen in bad condition was found in Meudon (Seine-et-Oise). Like Rageau *et al.* (1970) we consider that this record is doubtful as it is well outside the known range of the species (Mediterranean basin, especially North Africa). A second report (Ségui, 1966) mentions three discoveries of the species (124 larvae) in Corsica in 1966. Even though this report seems more reasonable from the biogeographical point of view, no specimens were preserved and the taxon is therefore excluded from the list until further surveys provide confirmation.

Aedes (Ochlerotatus) nigripes (Zetterstedt 1938) was mentioned by Séguy (1923) (4 females from Fontainebleau, June 1869, Dufour's collection) and quoted by Hedeën (1959) on the same basis. On the other hand, Callot & Rioux (1965) excluded it from their list. Considering the extremely northerly circumpolar range of the species, it is most unlikely to belong to the French fauna.

Aedes (Ochlerotatus) caspius hargreavesi Edwards 1920 was described from Italy and is mentioned as present in France in the catalogues of Knight & Stone (1977) and Dahl & White (1978). Séguy (1923) considered that the species could possibly occur in the south of France. Lack of precise records and a general absence of data regarding this sub-species, the validity of which requires confirmation, prevents us from retaining it in our lists.

Culex (Maillotia) deserticola Kirkpatrick 1924 was recorded in Corsica by Séguy (1965) who stated that he had collected the species in the larval stage on five occasions during his surveys of that year. He also recorded larvae of *Cx. hortensis*. Differences between larvae of these two species are extremely small. For reasons of biogeographic disjunction (the species is known from North Africa and Asia Minor (Dahl & White, 1978)), and the absence of preserved specimen, we do not consider the species as part of the fauna of France.

Culex (Culex) prosecutor Séguy 1927 (= *Culex pseudomimeticus* Séguy 1925) was described by Séguy (1925b, 1927) from an incomplete larva found in the Camargue. Mattingly considers it to be a synonym of *Cx. impudicus* (in Senevet & Andarelli, 1959). Although it is mentioned as present in France in various catalogues (Knight & Stone, 1977; Dahl & White, 1978; Minar, 1990), no new information has been published regarding this taxon. The description of the extremity of the abdomen matches that of larvae of the sub-genus *Neoculex*, which is represented in France (and in Europe) by *Cx. impudicus*, *Cx. martinii* and *Cx. territans*. No major taxonomic characters allow differentiation of the larvae of these three species. Both *Cx. impudicus* and *Cx. martinii* can be encountered in the area where Séguy found the larvae. Because of the lack of information about this specimen, we are unable to attribute it to any of these species. Therefore, we propose classifying *Cx. prosecutor* as *nomina dubia*.

IV. Species that could be encountered in France

We consider that the following species have a reasonable chance of being encountered in France (including Corsica):

Anopheles (Cellia) cinereus hispaniola (Theobald 1903): this species is common in Sardinia, and could therefore be discovered in Corsica (Aitken, 1954; Rageau *et al.*, 1970).

Aedes (Ochlerotatus) leucomelas (Meigen 1804) and *Culiseta (Culicella) ochroptera* (Peus 1935): both species occur in Germany, close to the French border (Oberrheinebene and Schwarzwald respectively) (Becker & Ludwig, 1981). Populations could perhaps be discovered in the north-east of France.

Aedes (Stegomyia) albopictus (Skuse 1894) and *Aedes (Ochlerotatus) atropalpus* s.s. (Coquillett 1902): these species were introduced into Italy with used tyres imported from North America. The former occurs in Piedmont and Sardinia; it first appeared in 1990 and has ever since expanded its range (Knudsen *et al.*, 1996). The latter was discovered in 1996 in Venice (Romi *et al.*, 1997). *Ae. albopictus* poses health threats, due to its high capacity to transmit arboviruses. France is considered a "high introduction risk" country (Knudsen *et al.*, 1996), and it is important to monitor closely the potential breeding niches (tyres, various containers, hollow trees) in the south of France and Corsica.

Aedes (Finlaya) echinus (Edwards 1920), *Culex (Culex) laticinctus* Edwards 1913 and *Culex (Culex) univittatus* Theobald 1901: as mentioned earlier, these Mediterranean species have not been positively recorded in France, but could nevertheless be present in the south of France and, in particular, Corsica.

V. Some features of the distribution and ecology of the mosquitoes of France

- *Species occurring in Corsica only: *An. marteri*, *An. labranchiae*, *An. sacharovi*, *An. superpictus*, *Cx. brumpti*.
- *Species confined to the south of France: *An. petraganaii*, *An. melanoon*, *An. subalpinus*, *An. hyrcanus*, *Ae. vittatus*, *Ae. berlandi*, *Ae. mariae*, *Ae. pulcritarsis pulcritarsis*, *Cx. mimeticus*, *Ur. unguiculata*.
- *Species limited to the north-east of France: *Ae. esoensis rossicus*, *Ae. dantaesus*, *Ae. nigrinus*, *Cs. alaskaensis*, *Cs. glaphyoptera*.
- *Mountain species: *Ae. pullatus*, *Ae. surcoufi*, *Ae. cataphylla*, *Cs. glaphyoptera*.
- *Saltwater species: *Ae. detritus*, *Ae. caspius caspius*, *Ae. dorsalis*.
- *Treehole species: *An. plumbeus*, *Ae. geniculatus*, *Ae. berlandi*, *Ae. pulcritarsis pulcritarsis*, *Or. pulcripalpis*.
- *Rare species: *Ae. vittatus*, *Ae. dantaesus*, *Ae. nigrinus*, *Ae. flavescens*, *Ae. refiki*, *Cx. brumpti*, *Cx. martinii*.

Conclusion

The French *Culicidae* fauna comprises 63 species to date. Even though this fauna is relatively well known, this list is not definitive. Indeed, thorough investigations of poorly studied areas and border regions could result in the addition of many species currently recorded in neighbouring countries. Additionally, global climatic change will very probably result in a northward extension of the range of some species. Systematic monitoring would allow the detection of such new arrivals, whether due to range extensions or accidental introductions.

Acknowledgements — This paper was written within the scientific programme of the *limitation de la nuisance due aux moustiques* which is funded by the Conseil Général du Haut-Rhin. I also wish to thank my friend Jacques Brunhes (ORSTOM) for advice on presentation. I also wish to thank the editors for translating the paper into English.

REFERENCES

- Agoulon, A. (1996) *Écologie de deux Aedes halophiles du littoral atlantique français: Aedes (Ochlerotatus) detritus* (Haliday, 1833) et *Aedes (Ochlerotatus) caspius* (Pallas, 1771) (Diptera: Culicidae). *Identification génétique des populations. Recherche d'autogénèse*. Thèse Universitaire Nantes, 180pp + annexes.
- Aitken, T.G.H. (1954) The Culicidae of Sardinia and Corsica (Diptera). *Bulletin of Entomological Research* 45, 437-494.
- Arnaud, J.D., Rioux, J.-A., Croset, H. & Guilvard, E. (1976) *Aedes (Ochlerotatus) surcoufi* (Theobald, 1912). Rétablissement du binôme ; analyse morphologique ; position au sein du complexe holarctique "excrucians". *Annales de Parasitologie* 51, 477-494.
- Becker, N. & Ludwig, H.W. (1981) Untersuchung zur Faunistik und Ökologie der Stechmücken (Culicinae) und ihrer Pathogene im Oberrheingebiet. *Mitteilung der deutsche Gesellschaft für allgemeine angewandte Entomologie* 2, 186-194.
- Brutus, L., Riandey, M.-F., Guilloteau, J., Monteny, N., Sannier, C. & Marjolet, M. (1994) Mise en évidence des deux espèces jumelles A et B du complexe *Aedes detritus* (Haliday, 1833) sur le littoral atlantique français. *Parasite* 1, 167-170.
- Bullini, L., Bianchi Bullini, A., Cianchi, R., Sabatini, A. & Coluzzi, M. (1980) Tassonomia biochemica del complesso *Anopheles maculipennis*. *Parasitologia* 22, 290-292.
- Callot, J. & Rioux J.-A. (1965) Liste sommaire des Culicidés de France. *Annales de Parasitologie humaine et comparée* 40, 242-245.
- Cianchi, R., Urbanelli, A., Sabatini, A., Coluzzi, M., Tordi, M.P. & Bullini, L. (1987) Electrophoretic evidence of reproductive isolation between sympatric populations of *Anopheles melanoon* and *An. subalpinus*. In: 3rd International Congress on Malaria and Babesiosis, p1560. International Laveran Foundation, 7-11 September 1987. Annecy, France.
- Cianchi, R., Sabatini, A., Boccolini, D., Bullini, L. & Coluzzi, M. (1980) Due entità riproduttivamente isolate sotto il nome di *Aedes caspius* (Diptera, Culicidae). *Atti XII Congresso Nazionale Italiano Entomologia, Roma* 2, 269-272.
- Dahl, C. & White, G.B. (1978) Culicidae. In *Limnofauna Europaea*, Illies ed., G. Fischer Verlag, Stuttgart, 390-396.
- Edwards, F.W. (1928) The Nematoceros Diptera of Corsica. *Diptera* 4, 157-189.

- Encinas Grandes, A. (1982) *Taxonomia y biologia de los mosquitos del area Salmantina (Diptera, Culicidae)*. Universidad de Salamanca. 437 pp.
- Gaffigan, T.V. & Ward, R.A. (1985) Index to the Second Supplement to *A catalog of the mosquitoes of the World*, with corrections and additions (Diptera: Culicidae). *Mosquito Systematics* 17, 52-63.
- Galliard, H. (1927) Notes sur les Culicines de Corse. *Annales de Parasitologie humaine et comparée* 5, 97-104.
- Galliard, H. (1928) Quelques Culicidés nouveaux pour la Corse en particulier *Culex impudicus* Ficalbi. *Annales de Parasitologie humaine et comparée* 6, 451-454.
- Galliard, H. (1931) *Culex brumpti* n. sp. moustique nouveau trouvé en Corse. *Annales de Parasitologie humaine et comparée* 9, 134-139.
- Harbach, R.E. (1994) Review of the internal classification of the genus *Anopheles* (Diptera: Culicidae): the foundation for comparative systematics and phylogenetic research. *Bulletin of Entomological Research* 84, 331-342.
- Hedeen, R.-A. (1958) A review of the mosquito larvae of France. I. Genera *Culiseta*, *Mansonia*, *Orthopodomyia*, and *Uranotaenia*. *Mosquito News* 18, 308-321.
- Hedeen, R.A. (1959) A review of the mosquito larvae of France. II. The genus *Aedes*. *Mosquito News* 19, 179-183.
- Knight, K.L. (1978) Supplement to *A catalog of the mosquitoes of the world* (Diptera: Culicidae). Thomas Say Foundation, Supp. 6, 107 pp.
- Knight, K.L. & Stone, A. (1977) *A catalog of the mosquitoes of the world* (Diptera: Culicidae). 2nd ed., Thomas Say Foundation, 6, 611 pp.
- Knudsen, A.B., Romi, R. & Majori, G. (1996) Occurrence and spread in Italy of *Aedes albopictus*, with implications for its introduction into other parts of Europe. *Journal of the American Mosquito Control Association* 12, 177-183.
- Lanzalavi, P. (1965) *Recherches sur les Culicidae (Nematocera) de la Corse*. Marseille, thèse Médecine, 97 pp.
- Minar, J. (1990) Family Culicidae in *Catalogue of Palaearctic Diptera*. Vol. 2. Soos & Papp ed. Budapest, 499 pp. 74-113.
- Mohrig, W. (1969) *Die Culiciden Deutschlands, Untersuchung zur Taxonomie, Biologie und Ökologie der einheimischen Stechmücken*. Parasitologische Schriftenreihe, G. Fischer Verlag, Jena, 18, 260 pp.
- Moussiegt, O. (1986) *Moustiques de France. Bibliographie et répartition. Inventaires Faune et Flore n°30*. Museum National D'Histoire Naturelle. Paris, 184 pp.
- Pasteur, N., Rioux, J.-A., Guilvard, E., Pech-Perieres, M.-J. & Verdier, J.-M. (1977) Existence chez *Aedes (Ochlerotatus) detritus* (Haliday, 1833) (Diptera-Culicidae) de Camargue de deux formes sympatriques et sexuellement isolées (espèces jumelles). *Annales de Parasitologie* 52, 325-337.
- Peus, F. (1970) Bemerkenswerte Mücken am Tegeler Fliess. *Berliner Naturschutzblätter*. Special number (May), 18-26.
- Rageau, J., Mouchet, J. & Abonnec, E. (1970) Répartition géographique des moustiques (Diptera: Culicidae) en France. *Cahiers O.R.S.T.O.M., série Entomologie médicale et Parasitologie* 8, 289-315.
- Ribeiro, H., Da Cunha Ramos, H., Pires, C.A. & Antunes Capela, R. (1988) An annotated checklist of the mosquitoes of continental Portugal (Diptera: Culicidae). *Actas III Congreso Iberico de Entomologia Granada*, 233-253.
- Rioux, J.-A. & Ruffié, J. (1957) Présence d'*Anopheles maculipennis* s.sp. *subalpinus* Hackett et Lewis 1937, en Bas-Languedoc. Intérêt des méthodes caryosystématiques et statistiques. *Bulletin de la Société de Pathologie exotique* 50, 831-837.
- Roman, E. (1944) Un *Aedes* nouveau pour la France. Localités inédites de Moustiques peu répandus. *Bulletin de la Société entomologique de France* 49, 35-36.
- Roman, E. (1958) Contribution à la répartition en France des diptères de la famille des Culicidae. Gîtes larvaires particuliers de quelques-uns d'entre eux. *Annales de Parasitologie* 33, 115-130.
- Roman, E. & Morel, P. (1946) Moustiques de l'Ouest Savoyard. *Bulletin de la Société de Pathologie exotique* 29, 372-376.
- Roman, E. & Pichot, J. (1976) Diptères Culicidae de la faune lyonnaise : moustiques nouveaux, renseignements complémentaires sur certains déjà signalés. *Bulletin de la Société Linnéenne de Lyon* 45, 57-60.
- Romi, R., Sabatinelli, G., Giannuzzi Savelli, L., Raris, M., Zago, M. & Malatesta, R. (1997) Identification of a north American Mosquito Species, *Aedes atropalpus* (Diptera: Culicidae), in Italy. *Journal of the American Mosquito Control Association* 13, 245-246.
- Salieres, A., Guy, Y., Suzzoni-Blatger, J. & Cousserans, J. (1978) Bilan de quatre années de recherches sur le "complexe *maculipennis*" (Diptera - Culicidae- Anophelinae). *Annales de Parasitologie* 53, 751-756.

- Schaffner, F. (1992) *Les Moustiques de Haute-Alsace. I - Systématique*. Retirage de thèse D.R.S., Conseil Général du Haut-Rhin, 295 pp.
- Schaffner, F. & Pfirsch, F. (1995) Un nouveau moustique pour la faune française: *Aedes (Ae.) esoensis rossicus* Dolbeskin, Gorickaja & Mitrofanova, 1930 (Diptera, Culicidae). *Bulletin de la Société entomologique de France* 100, 325-330.
- Séguy, V. (1965) *Enquête entomologique. Année 1965*. Rapport Service Antipaludique de la Corse. D.A.S.S., 37 pp.
- Séguy, V. (1966) *Enquête entomologique n°3*. Rapport Service Antipaludique de la Corse. D.A.S.S., 66 pp.
- Séguy, E. (1920a) Notes sur quelques moustiques (Dipt.) peu connus ou nouveaux pour la faune française. *Bulletin de la Société entomologique de France* 15, 251-25.
- Séguy, E. (1920b) Les moustiques de France. *Culicinae*. *Bulletin du Muséum National d'Histoire Naturelle* 26, 322-329.
- Séguy, E. (1921) Les moustiques de France. *Culicinae*. *Bulletin du Muséum National d'Histoire Naturelle* 27, 162-169.
- Séguy, E. (1923) *Les moustiques de France. (Histoire naturelle des moustiques de France. Etude systématique et biologique des moustiques de l'Europe centrale et septentrionale et de leurs parasites)*. Lechevalier, Paris, 225 pp.
- Séguy, E. (1925a) *Faune de France. 12. Diptères (Nématocères piqueurs)*. Lechevalier, Paris, 109 pp.
- Séguy, E. (1925b) Notes sur les moustiques de l'Afrique Mineure, de l'Égypte et de la Syrie. I. *Encyclopédie Entomologique, série B, Diptera* 2, 13-21.
- Séguy, E. (1927) Notes sur les moustiques de l'Afrique Mineure, de l'Égypte et de la Syrie. II. *Encyclopédie Entomologique, série B, Diptera* 4, 27-28.
- Séguy, E. (1932) Notes sur les moustiques IV. *Encyclopédie Entomologique, série B, Diptera* 6, 97-102.
- Sénevet, G. & Andarelli, L. (1959) *Les moustiques de l'Afrique du Nord et du bassin méditerranéen. Les genres Culex, Uranotaenia, Theobaldia, Orthopodomyia et Mansonia*. Lechevalier, Paris, 384 pp.
- Shevchenko, A.K. & Prudkina, N.S. (1973) On the morphology of the genitalia in mosquito males from the genus *Aedes*. *Vestnik Zoologii* 6, 40-47.
- Ward, R.A. (1984) Second Supplement to *A catalog of the mosquitoes of the world (Diptera: Culicidae)*. *Mosquito Systematics* 16, 227-270.
- Ward, R.A. (1992) Third Supplement to *A catalog of the mosquitoes of the world (Diptera: Culicidae)*. *Mosquito Systematics* 24, 177-230.

Culicidae Workshop - Fourth International Congress of Dipterology

Ralph Harbach and Keith Snow

Department of Entomology, The Natural History Museum, Cromwell Road, London SW7 5BD and Department of Environmental Sciences, University of East London, Romford Road, London E15 4LZ

We take this opportunity to remind culicidologists to participate in the Culicidae Workshop at the Fourth International Congress of Dipterology (6-13 September 1998, Keble College, Oxford, England) which will focus on 'Trends in Mosquito Ecology and Systematics.'

The keynote presentation: *Overview of the status and current advances in mosquito systematics* will be given by Tom Zavortink and there will be a full programme of both oral and poster presentations.